


FORM PTO-1390 U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE REV. 2/01T		ATTORNEY'S DOCKET NUMBER <b>04853.0089</b> Customer No.: 22,852
<b>TRANSMITTAL LETTER TO THE UNITED STATES          DESIGNATED/ELECTED OFFICE (DO/EO/US)          CONCERNING A FILING UNDER 35 U.S.C. 371</b>		U.S. APPLICATION NO. (If known, see 37CFR 1.5) <b>10/069251</b> ✓
INTERNATIONAL APPLICATION NO. <b>PCT/JP00/05686</b> ✓	INTERNATIONAL FILING DATE <b>August 24, 2000</b> ✓	PRIORITY DATE CLAIMED <b>August 24, 1999</b> ✓
TITLE OF INVENTION: <b>ANIMALS WITH KNOCKED OUT <math>\alpha</math>-TOCOPHEROL TRANSFER PROTEIN GENE</b>		
APPLICANTS FOR DO/EO/US: <b>1) Keizo INOUE, 2) Hiroyuki ARAI, 3) Makoto ARITA, 4) Kou-ichi JISHAGE, and 5) Hiroshi SUZUKI</b>		
Applicants herewith submit to the United States Designated/Elected Office (DO/EO/US) the following items and other information:		
1.	<input checked="" type="checkbox"/>	This is a <b>FIRST</b> submission of items concerning a filing under 35 U.S.C. 371.
2.	<input type="checkbox"/>	This is a <b>SECOND</b> or <b>SUBSEQUENT</b> submission of items concerning a filing under 35 U.S.C. 371.
3.	<input type="checkbox"/>	This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below.
4.	<input checked="" type="checkbox"/>	The US has been elected by the expiration of 19 months from the priority date (Article 31).
5.	<input checked="" type="checkbox"/>	A copy of the International Application as filed (35 U.S.C. 371 (c)(2)).
	a. <input type="checkbox"/>	is attached hereto (required only if not communicated by the International Bureau).
	b. <input checked="" type="checkbox"/>	has been communicated by the International Bureau.
	c. <input type="checkbox"/>	is not required, as the application was filed with the United States Receiving Office (RO/US).
6.	<input checked="" type="checkbox"/>	An English language translation of the International Application as filed (35 U.S.C. 371 (c)(2)).
	a. <input checked="" type="checkbox"/>	is attached hereto.
	b. <input type="checkbox"/>	has been previously submitted under 35 U.S.C. 154 (d)(4).
7.	<input checked="" type="checkbox"/>	Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3)).
	a. <input type="checkbox"/>	are attached hereto (required only if not communicated by the International Bureau).
	b. <input type="checkbox"/>	have been communicated by the International Bureau.
	c. <input type="checkbox"/>	have not been made; however, the time limit for making such amendments has NOT expired.
	d. <input checked="" type="checkbox"/>	have not been made and will not be made.
8.	<input type="checkbox"/>	An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).
9.	<input type="checkbox"/>	An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).
10.	<input type="checkbox"/>	An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).
<b>Items 11 to 20 below concern document(s) or information included:</b>		
11.	<input checked="" type="checkbox"/>	Information Disclosure Statement under 37 CFR 1.97 and 1.98
12.	<input type="checkbox"/>	An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13.	<input type="checkbox"/>	A FIRST preliminary amendment.
14.	<input type="checkbox"/>	A SECOND or SUBSEQUENT preliminary amendment.
15.	<input type="checkbox"/>	A Substitute specification.
16.	<input type="checkbox"/>	A change of power of attorney and/or address letter.
17.	<input type="checkbox"/>	A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821-1.825.
18.	<input type="checkbox"/>	A second copy of the published international application under 35 U.S.C. 154 (d)(4).
19.	<input type="checkbox"/>	A second copy of the English language translation of the international application 35 U.S.C. 154 (d)(4).
20.	<input checked="" type="checkbox"/>	Other items or information:
	a. <input checked="" type="checkbox"/>	Copy of cover page of International Publication No. WO 01/13716 A1.
	b. <input checked="" type="checkbox"/>	Statement.
	c. <input checked="" type="checkbox"/>	Sequence Listing - 12 pages
	d. <input checked="" type="checkbox"/>	Computer Readable Sequence Listing - One Diskette.

U.S. APPLICATION NO. (If known, see 37CFR 1.5)		INTERNATIONAL APPLICATION NO.		ATTORNEY'S DOCKET NUMBER	
10/069251		PCT/JP00/05686		04853.0089	
21. <input checked="" type="checkbox"/> The following fees are submitted:				CALCULATIONS PTO USE ONLY	
BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):					
Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO .....				\$1040.00	
International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO .....				\$890.00	
International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search fee (37 CFR 1.445(a)(2)) paid to USPTO .....				\$740.00	
International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) .....				\$710.00	
International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33 (1)-(4) .....				\$100.00	
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$890.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than months from the earliest claimed priority date (37 CFR 1.492 (e)).				<input type="checkbox"/> 20 <input type="checkbox"/> 30	\$
CLAIMS	NUMBER FILED		NUMBER EXTRA	RATE	
Total Claims	17	- 20 =	0	x \$18.00	\$
Independent Claims	2	- 3 =	0	x \$84.00	\$
<input checked="" type="checkbox"/> MULTIPLE DEPENDENT CLAIMS (if applicable)				+\$280.00	\$280.00
TOTAL OF THE ABOVE CALCULATIONS =				\$1170.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$	
SUBTOTAL =				\$1170.00	
Processing fee of \$130.00 for furnishing the English translation later than months from the earliest priority date (37 CFR 1.492(f)).				<input type="checkbox"/> 20 <input type="checkbox"/> 30	\$
TOTAL NATIONAL FEE =				\$1170.00	
Fee for recording the enclosed assignment (37 CFR 1.21 (h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property.				+	\$
TOTAL FEES ENCLOSED =				\$1170.00	
				Amount to be refunded:	\$
				charged:	\$
a. <input checked="" type="checkbox"/> A check in the amount of \$ 1170.00 to cover the above fees is enclosed.					
b. <input type="checkbox"/> Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed.					
c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 06-0916. A duplicate copy of this sheet is enclosed.					
d. <input type="checkbox"/> Fees are to be charged to a credit card. <b>WARNING:</b> Information on this form may become public. <b>Credit card information should not be included on this form.</b> Provide credit card information and authorization on PTO-2038.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137 (a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO:					
Finnegan, Henderson, Farabow, Garrett & Dunner, L.L.P.					
1300 I Street, N.W.					
Washington, D.C. 20005-3315					
EFC/FPD/sci					
DATED: February 22, 2002					
					
				SIGNATURE	
				Ernest F. Chapman/25,961	
				NAME/REGISTRATION NO.	

Attorney Docket No.: 04853.0089  
Customer Number 22,852

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: )  
)  
**Keizo INOUE** et al. )  
) Group Art Unit: Not Yet Assigned  
Serial No.: Not Yet Assigned )  
) Examiner: Not Yet Assigned  
Filed: February 22, 2002 )  
)  
National Stage of International )  
Application No. **PCT/JP00/05686**, under )  
35 U.S.C. 371, for **ANIMALS WITH** )  
**KNOCKED OUT  $\alpha$ -TOCOPHEROL** )  
**TRANSFER PROTEIN GENE** )  
)

Assistant Commissioner for Patents  
Washington, DC 20231

Sir:


STATEMENT

The information recorded in computer readable form (attached diskette) is  
identical to the written sequence listing..

Respectfully submitted,

FINNEGAN, HENDERSON, FARABOW,  
GARRETT & DUNNER, L.L.P.

Dated: February 22, 2002

By:   
Ernest F. Chapman  
Reg. No. 25,961

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

EFC/FPD/sci  
Enclosures

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

## DESCRIPTION

ANIMALS WITH KNOCKED OUT  $\alpha$ -TOCOPHEROL TRANSFER PROTEIN GENETECHNICAL FIELD OF THE INVENTION

The present invention relates to a mammal modified to inhibit the expression of its gene for  $\alpha$ -tocopherol transfer protein (hereinafter, referred to as " $\alpha$ -TTP"). Human  $\alpha$ -TTP gene is a causative gene for familial isolated vitamin E deficiency. The mammal can therefore be used for the development of a therapeutic method and/or agent for the disease, as well as for the development of a therapeutic method and/or agent for oxidative stress-induced diseases such as arteriosclerosis or diabetes.

BACKGROUND ART

Oxygen is essential for living organisms, but it also has a dangerous aspect of causing unwanted oxidation of organisms' components. Living organisms have a series of defense mechanisms against such an oxidative stress. Vitamin E is one of important anti-oxidative substances in the living organisms. Among organism's components, a biomembrane phospholipid (particularly, a polyunsaturated fatty acid chain in phospholipid) is most likely to receive an attack by oxygen. When biomembranes undergo an oxidative impairment, the membrane permeability increases, and finally leading to cell death. Vitamin E is lipid-soluble (lipophilic) and usually buried in biomembrane bilayers in cells. It plays the most dominant role in preventing biomembrane oxidation.

As a protein capable of specifically binding to this vitamin E,  $\alpha$ -TTP was isolated from a soluble fraction of rat liver and found to enhance intermembrane transfer (Eur. J. Biochem. 177, 537, 1981). In addition, purification and gene structure analysis were also performed on this protein (J. Biol. Chem. 268, 17705, 1993; Biochem. J. 306, 437, 1995). Further, human  $\alpha$ -TTP gene was shown to be a causative gene per se for a

hereditary disease called “familial isolated vitamin E deficiency (FIVE deficiency)” (Nature Genetics 9, 141, 1995).

Familial isolated vitamin E deficiency has previously been known as a hereditary disease which produces no increase in vitamin E level in the body, even when much vitamin E is taken into the body. A patient with this disease suffers from necrosis of nerves, particularly sensory nerves, and in a serious case he will die around 20 years of age. Vitamin E circulates in the blood as a conjugate with plasma lipoprotein and then enters peripheral tissues. Meanwhile, plasma lipoprotein is secreted from the liver and finally returns to the liver to be metabolized. Vitamin E absorbed from food through blood vessels is incorporated onto lipoprotein (VLDL) in the liver. In this hereditary disease, an impairment is found in just this process, i.e., an incorporation process of vitamin E onto VLDL in the liver.

#### PROBLEMS TO BE SOLVED BY THE INVENTION

In view of these circumstances, the object of the present invention is to provide a non-human mammal useful in analyzing  $\alpha$ -TTP functions and developing a therapeutic agent for diseases caused by  $\alpha$ -TTP mutation. In the prior art, an animal could enter a vitamin E deficient state by feeding it with a vitamin E-deficient diet, but in a normal animal, a very long period of time was required to achieve complete clearance of vitamin E from the animal body because the normal animal has an efficient re-circulation system for vitamin E (this system involves  $\alpha$ -TTP). In the present invention, an animal with a disrupted  $\alpha$ -TTP gene involved in this vitamin E re-circulation system is created to provide a congenitally vitamin E-deficient animal model highly sensitive to oxidative stress, which is advantageous in understanding diseases caused by vitamin E deficiency, i.e., deficiency of anti-oxidative substances critical for the body, and in developing a therapeutic agent for such diseases. More specifically, the object of the present invention is to provide a knockout animal artificially modified to inhibit  $\alpha$ -TTP gene expression and a method for producing the

animal.

### MEANS FOR SOLVING THE PROBLEMS

We tried to generate a mammal model with an artificially disrupted  $\alpha$ -TTP gene. Specifically, as shown in the Examples in more detail, mouse  $\alpha$ -TTP gene (cDNA, genomic DNA) was cloned and used to construct a vector for homologous recombination, which was then introduced into mouse embryonic stem cells (ES cells) to obtain a recombinant clone. The recombinant clone was then transferred to a recipient mouse, thereby successfully obtaining a mouse with a mutated  $\alpha$ -TTP gene. Mammals obtained according to the present invention or established cell lines prepared therefrom are considered to be useful tools for understanding mechanisms for the development of various diseases caused by a disrupted  $\alpha$ -TTP gene and other disease conditions suspected to be associated with oxidative stress (e.g., arteriosclerosis, diabetes, ischemic diseases, Parkinson's disease) and further considered to be very useful tools for developing a therapeutic method and/or agent for these diseases. They are therefore expected to be used for a variety of purposes.

The present invention relates to the following embodiments (1) to (4):

- (1) a non-human mammal artificially modified to inhibit the expression of its endogenous gene encoding  $\alpha$ -TTP, and a non-human mammal cell prepared therefrom;
- (2) a non-human mammal cell artificially modified to inhibit the expression of its endogenous gene encoding  $\alpha$ -TTP, said cell having the ability to differentiate into an individual;
- (3) a method for producing the non-human mammal of (1) above, which comprises the steps of: (a) inserting the non-human mammal cell of (2) above into an embryo taken from a pregnant female, and (b) transferring the embryo into the uterus of a pseudopregnant female; and
- (4) a method for screening medicaments, which comprises using the non-human mammal of (1) above or the non-human mammal cell of (1) or (2) above, and a

medicament obtained thereby.

#### DISCLOSURE OF THE INVENTION

The present invention will be described below in more detail.

##### (1) Knockout animals and cells prepared therefrom

The knockout animal and cells prepared therefrom of the present invention are characterized by being artificially modified to inhibit the expression of their endogenous gene encoding  $\alpha$ -TTP.

Examples of a means for modifying an animal or cell to inhibit gene expression include, but are not limited to, a means for disrupting a part of the  $\alpha$ -TTP gene or an expression regulatory region thereof. As used herein, the phrase “inhibit gene expression” or “inhibit the expression” encompasses both complete and partial inhibition. It also encompasses inhibition caused in a particular environment. Further, it encompasses a case where one of two alleles is inhibited from expressing.

Any type of animal may be used, so long as it is a mammal other than human. Preferred animals are those belonging to Rodent, including mouse, rat, hamster and rabbit, with mouse being particularly preferred among these.

The knockout animal of the present invention may be prepared, for example, in the manner stated below.

The knockout animal of the present invention is useful in developing a therapeutic agent and/or method for diseases caused by impaired functions of the  $\alpha$ -TTP gene and/or diseases induced by oxidative stress. For example, the knockout animal of the present invention is administered with a test compound to assay the compound for its effect on arteriosclerosis or diabetes, thereby selecting a compound having a desired

effect. This enables the acquisition of a potential candidate for a therapeutic agent.

In addition, cells prepared from the knockout animal of the present invention are contemplated to be used for the development of a therapeutic agent and/or method for the diseases mentioned above. For example, cells are prepared from an embryo or the like of the knockout animal according to the present invention, and then contacted with a test compound to assay the compound for its effect on oxidative impairment of the cell membranes etc., thereby selecting a compound having a desired effect. Cells to be used may be either primary cultured cells or established cell lines prepared therefrom. The compound thus screened may be a potential candidate for a medicament.

## (2) Animal cells

The animal cells of the present invention are characterized by being artificially modified to inhibit the expression of their endogenous gene encoding  $\alpha$ -TTP and by their ability to differentiate into an individual.

A means for modifying a cell to inhibit gene expression and the meaning thereof, as well as the type of animal to be used are as defined for the above knockout animal.

The animal cells of the present invention may be prepared in any manner. To prepare cells modified to inhibit  $\alpha$ -TTP gene expression through disruption of at least a part of the  $\alpha$ -TTP gene or an expression regulatory region thereof, a homologous recombination (knockout) vector may be constructed and introduced into appropriate cells.

The homologous recombination vector comprises a nucleotide sequence designed to inactivate an endogenous  $\alpha$ -TTP gene of a target animal. Such a



nucleotide sequence may be, for example, a nucleotide sequence lacking at least a part of the  $\alpha$ -TTP gene or an expression regulatory region thereof or a nucleotide sequence containing another gene inserted into the sequence of the  $\alpha$ -TTP gene or an expression regulatory region thereof. Preferably, another gene thus inserted into the sequence of the  $\alpha$ -TTP gene or an expression regulatory region thereof may also function as a marker. Examples of such a gene include a drug-resistance gene such as neomycin-resistance gene (for G418 resistance selection) and thymidine kinase gene (for ganciclovir resistance selection); a toxin gene such as diphtheria toxin (DT) A gene; or combinations thereof. These genes may be inserted into any site on the  $\alpha$ -TTP gene, so long as the inserted gene can inhibit the expression of an endogenous  $\alpha$ -TTP gene in a target.

The insertion of these genes into the sequence of the cloned  $\alpha$ -TTP gene may be accomplished *in vitro* by standard DNA recombination techniques (See, Sambrook, J. et al., Molecular Cloning, Cold Spring Harbor Laboratory Press (1989)).

The homologous recombination vector thus constructed is introduced into cells having the ability to differentiate into an individual (e.g., ES cells) to cause homologous recombination between the vector and the  $\alpha$ -TTP gene in the cells.

The homologous recombination vector may be introduced into the cells in a manner well known to those skilled in the art, for example, by using an electroporation technique. As a result, in some of the cells, homologous recombination occurs between the cellular  $\alpha$ -TTP gene and the corresponding region of the homologous recombination vector, thereby replacing the wild-type gene by a gene constructed in the homologous recombination vector. Thus, cells having a modified  $\alpha$ -TTP gene containing the inserted marker gene can be obtained.

In a case where a marker gene is used in the homologous recombination vector,

cells undergoing a desired homologous recombination event will have an inactivated  $\alpha$ -TTP gene and will also attain the marker gene, thereby allowing the use of this marker gene as an indicator for cell screening. For example, when a drug resistance gene is used as a marker gene, cells which have been subjected to vector introduction may be screened for a desired homologous recombination event by culturing the cells in the presence of the drug at a lethal level.

The animal cells of the present invention may be used for production of the knockout animal stated above. These cells may also be used for screening of medicaments, as in the case of the cells prepared from the knockout animal according to the present invention.

### (3) Method for production of non-human mammals

The method for producing a non-human mammal of the present invention comprises the steps of: (a) inserting the non-human mammal cell described above into an embryo taken from a pregnant female to form a chimeric embryo, and (b) transferring the chimeric embryo into the uterus of a pseudopregnant female.

#### (a) Step of forming chimeric embryos

In a case where ES cells are used for insertion into embryos, these cells may be injected into blastocysts to form chimeric embryos. Blastocysts to be used for injection may be obtained by flushing the uteri of pregnant females.

#### (b) Step of transferring chimeric embryos

The chimeric embryos may be transferred into the uterine horns of pseudopregnant mammals to obtain chimeric animals. In order to permit a determination in the resulting animals as to whether the injected cells (ES cells) have been successfully incorporated into developing embryos, it is desirable to choose blastocysts such that the resulting animals have a visual difference (e.g., coat color)

between parts originating from the injected cells and parts originating from blastocysts.

Following the above two steps, the resulting chimeric animal is crossed with an appropriate animal line of the same species to obtain pups. If the chimeric animal has germ cells derived from the injected cells, pups modified to inhibit  $\alpha$ -TTP gene expression can be obtained.

#### (4) Method for screening medicaments and medicaments obtained thereby

The method for screening medicaments of the present invention is characterized by using the non-human mammals or non-human mammal cells mentioned above.

The non-human mammal etc. of the present invention may be a model for diseases caused by impaired functions of the  $\alpha$ -TTP gene (e.g., familial isolated vitamin E deficiency) and diseases induced by oxidative stress (e.g., arteriosclerosis, diabetes, ischemic diseases, Parkinson's disease), and hence may be used to screen therapeutic and prophylactic agents for these diseases.

### EXAMPLES

The present invention will be further described in the following examples. The examples are provided for illustrative purposes only, and are not intended to limit the scope of the invention.

#### Example 1: Construction of homologous recombination vectors for $\alpha$ -TTP gene

In this example, to construct a homologous recombination vector for mouse  $\alpha$ -TTP gene, mouse genomic  $\alpha$ -TTP gene was first cloned. This genomic DNA was then used to construct a homologous recombination vector carrying both a neomycin-resistance gene and a thymidine kinase gene inserted thereinto for positive/negative selection, according to the report of Mansour et al. (Nature 336, 348,

1988). The neomycin-resistance gene was inserted to replace Exon 1 of  $\alpha$ -TTP gene such that no normal  $\alpha$ -TTP would be produced. Details will be described below.

#### A. Cloning of mouse $\alpha$ -TTP gene

A C57BL/6 mouse liver cDNA library (Lambda gt22A) was prepared using a SUPERScript Lambda System (GIBCO BRL). Plaque hybridization was performed on this library using the entire open reading frame of rat TTP cDNA as a probe to obtain two positive clones, neither of which extended to the translation initiation site. In turn, RT-PCR was performed on total RNA prepared from mouse liver using MATTP-00 (AGGAATTCATGGCAGAGATGCG; SEQ ID NO: 8) and MATTP-04 (AGGGCGTAGATCTGCACTTAAT; SEQ ID NO: 9) as primers. The sequence of MATTP-00 was obtained from a sequence of mouse genomic TTP DNA cloned separately. Sequencing for amplified products was performed to determine the entire open reading frame sequence of mouse TTP cDNA. The sequence of mouse TTP cDNA thus obtained is shown in SEQ ID NO: 1 and the amino acid sequence thereof is shown in SEQ ID NO: 2.

Mouse genomic  $\alpha$ -TTP DNA was cloned as follows.

A 129/SVJ mouse genomic DNA Lambda FIX II library was screened by plaque hybridization using the entire open reading frame of rat TTP cDNA as a probe. Phage DNA prepared from the resulting positive clone (BS-MG6) was cleaved with a restriction enzyme EcoRI and subjected to Southern blotting by using a 5'-terminal 260 bp sequence of rat TTP as a probe to detect a fragment containing an exon with the translation initiation site, which was then subcloned into pBluescriptII (TTP-MG6).

The nucleotide sequence of TTP-MG6 (2749 bp) was sequenced to confirm that Exon 1 was present in TTP-MG6 (SEQ ID NO: 3).

The clone BS-MG6 was cleaved with major restriction enzymes (EcoRI, SmaI, KpnI, ApaI, EcoRV, Sall, HindIII) to prepare a map showing the locations of cleavage sites for these restriction enzymes (Fig. 1).

#### B. Construction of homologous recombination vectors

The mouse  $\alpha$ -TTP gene contained in the clone BS-MG6 phage DNA was cleaved at EcoRI sites to give two fragments (Fragments 1 and 2; Fig. 1), each of which was then subcloned into a plasmid vector. More specifically, BS-MG6 phage DNA was first cleaved with EcoRI, and the resulting fragments of approximately 3.75 kbp (Fragment 1) and of approximately 5.5 kbp (Fragment 2) were then inserted into an EcoRI site of pBluescriptII (Toyobo), which had been modified to replace a SmaI site by a Sall site via introduction of a Sall linker (Takara), to obtain clones pSKot-2 and pSKot-3, respectively.

These subclones were used to construct a homologous recombination vector as follows. First, the clone pSKot-2 was cleaved with EcoRI/EcoRV and then blunt-ended by a technique using a Takara DNA Blunting Kit (Takara) to obtain pSKot-2-1. Subsequently, pSKot-2-1 was cleaved with SmaI, into which a HindIII linker was inserted (pSKot-2-2). A neomycin-resistance gene without a polyadenylation signal was integrated into a HindIII site of pSKot-2-2 to replace Exon 1 of  $\alpha$ -TTP by the neomycin-resistance gene (pSKot-2-2Neo). A BamHI Stop Codon linker (Nippon Gene Co., Ltd.) was then introduced into a SmaI site of Exon 2 in pSKot-3 (pSKot-3 stop codon). pSKot-2-2Neo was cleaved with EcoRI and then ligated to an EcoRI fragment cleaved from pSKot-3 stop codon (pSKot-2+3Neo). pSKot-2+3Neo was cleaved with XhoI and ligated to a thymidine kinase gene to obtain a homologous recombination vector ( $\alpha$ -TTP Targeting Vector).

The vector thus obtained is characterized by the following features (Fig. 2):

- (i) having a neomycin-resistance gene inserted to replace Exon 1 and a stop codon

sequence inserted into Exon 2;

- (ii) having a thymidine kinase gene as a marker gene for negative selection; and
- (iii) having regions homologous to the wild-type  $\alpha$ -TTP gene, in which one region of approx. 0.8 kb is upstream of, and the other of approx. 8 kb is downstream of the neomycin-resistance gene.

Example 2: Establishment of ES cells with mutated  $\alpha$ -TTP gene by homologous recombination

In this example, a homologous recombination vector was introduced into mouse ES cells (AB2.2-Prime ES Cells; The Mouse Kit, LEXICON) by electroporation and the cells were then selectively cultured in the presence of G418. The resulting G418-resistant colonies were assayed for homologous recombination events by PCR and Southern blotting. Details will be described below.

The homologous recombination vector ( $\alpha$ -TTP Targeting Vector) DNA (30  $\mu$ g) was cleaved with NotI to give linearized DNA, which was then purified. This DNA was suspended in electroporation buffer (ESQ PBS; The Mouse Kit, LEXICON) containing  $3 \times 10^7$  mouse ES cells (AB2.2-Prime ES Cells; The Mouse Kit, LEXICON) and subjected to gene transfer under the following conditions: a field strength of 575V/cm and a capacitance of 500  $\mu$ F. Twenty-four hours after the gene transfer, the cells were selectively cultured in the presence of G418 (Genetisin, Sigma) at a final concentration of 300  $\mu$ g/ml.

To culture the ES cells, a medium for ES cells (hereinafter, referred to as "ES Cell Medium") was prepared from ESQ DMEM medium (The Mouse Kit, LEXICON), which was supplemented with fetal bovine serum (ESQ FBS; The Mouse Kit, LEXICON) at a final concentration of 15%, L-glutamine (ESQ GPS; The Mouse Kit, LEXICON) at a final concentration of 2 mM,  $\beta$ -mercaptoethanol (ESQ BME; The Mouse Kit, LEXICON) at a final concentration of 100  $\mu$ M, penicillin at a final

concentration of 50 U/ml and streptomycin at a final concentration of 50 µg/ml.

In addition, ESQ Feeder cells (The Mouse Kit, LEXICON) were used as feeder cells for ES cells. To culture these feeder cells, ESQ DMEM medium supplemented with FBS at a final concentration of 7% was prepared. ESQ Feeder cells ( $5 \times 10^7$  cells/vial) were rapidly thawed at 37°C and then adjusted to a cell density of  $4.4 \times 10^5$  cells/ml with the medium for feeder cells. The resulting cell suspension was aliquoted into culture devices pre-coated with gelatin (ESQ Gelatin; The Mouse Kit, LEXICON) at an amount of 12 ml for 100 mm  $\phi$  dish, 4 ml for 60 mm  $\phi$  dish, 2 ml/well for 6-well plate, 0.5 ml/well for 24-well plate, and 75 µl/well for 96-well plate. The feeder cells thus prepared were used within 3 weeks.

Eleven days after the gene transfer, the appearing G418-resistant colonies were subcultured to a 96-well microplate as follows. Namely, each G418-resistant colony was transferred into 30 µl of ESQ trypsin solution (The Mouse Kit, LEXICON) per well of a 96-well microplate (Corning 2586OMP) by using a micropipette, and then treated for several minutes, followed by addition of ES Cell Culture Medium (70 µl), to prepare a single cell suspension by pipetting. Each of the resulting cell suspensions was transferred to another 96-well microplate (Falcon 3072) and further cultured. After 3 days, the cells grown to confluency on the 96-well microplate were divided into two groups as follows. Namely, the cells were dispersed in 25 µl of TE, followed by addition of ES Cell Medium (25 µl), to prepare a single cell suspension by pipetting. 50 µl of 2 × Freezing medium (ESQ DMEM:ESQ FBS:DMSO = 2:2:1; The Mouse Kit, LEXICON) was added to the suspension, 20 µl of which was then subcultured in 150 µl of ES Cell Medium per well of a gelatin-coated 96-well microplate (Iwaki 4860-020) and further cultured in order to extract DNA for use in PCR screening for homologous recombination events. The remaining ES cells were frozen and stored at -80°C with addition of liquid paraffin (100 µl; filter-sterilized through 0.2 µm filter). The ES cells used for DNA extraction were cultured in the absence of the feeder cells, whereas all ES

cells used for other purposes were cultured in the presence of the feeder cells. PCR screening for homologous recombination events was performed in the following manner. Namely, the medium was removed from each well of the 96-well plate containing the cells grown to confluency. Lysis buffer (5  $\mu$ l of 10  $\times$  Taq buffer, 5  $\mu$ l of 5% NP-40, 4  $\mu$ l of Proteinase K, 36  $\mu$ l of H<sub>2</sub>O) was added to each well and the plate was heated at 55°C for 2 hours. Each lysate sample was collected in a 0.5 ml tube and treated at 95°C for 15 minutes, followed by centrifugation at 10,000 rpm for 10 to 15 minutes. An aliquot (1  $\mu$ l) of each resulting supernatant was used as template DNA for PCR.

PCR primers were designed to amplify a region of approximately 0.9 kb located between PGK promoter of the neomycin-resistance gene on the homologous recombination vector and a region upstream of Exon 1 absent in the homologous recombination vector (Fig. 2).

More specifically, PCR was performed using PGK-1 primer containing a sequence on PGK promoter (5' GCTAAAGCGCATGCTCCAGACTGCCTTG 3'; SEQ ID NO: 5) and ot-198 primer located upstream of Exon 1 (5' AGCCCACACAAAAATGAAAAACGTCTCCAAG 3'; SEQ ID NO: 6) under the following conditions.

Reaction cocktail composition:

10  $\times$  ExTaq buffer (TaKaRa) 5  $\mu$ l  
 2.5 mM dNTPs 4  $\mu$ l  
 ExTaq (TaKaRa) 0.5  $\mu$ l  
 10  $\mu$ M ot-198 primer 1  $\mu$ l  
 10  $\mu$ M PGK-1 primer 1  $\mu$ l  
 Sample 1  $\mu$ l  
 H<sub>2</sub>O 37.5  $\mu$ l



Reaction conditions:

95°C for 1 minute → (94°C for 30 seconds → 62°C for 1 minute → 72°C for 1 minute and 20 seconds) × 35 cycles → 72°C for 7 minutes

Among 316 G418-resistant clones screened by PCR, 5 clones were identified as homologous recombinants (Clone L44, Clone L228, Clone L236, Clone L253 and Clone L254).

The clones, which had been confirmed to have undergone homologous recombination events by PCR screening, were subcultured to a 24-well plate from the freeze-stored 96-well plate after thawing at 37°C. This 24-well plate was cultured at 37°C for 24 hours and the medium was then replaced to remove DMSO and liquid paraffin. When the respective clones were grown to 75 to 90% confluency, they were subcultured to a 6-well plate from the 24-well plate. When two wells in the 6-well plate showed cell growth (75 to 90% confluency), one of these two wells was provided for freeze storage and the other was provided for injection into blastocysts and DNA extraction.

Freeze storage was performed as follows. Namely, the cells were rinsed twice with ESQ PBS, and then treated with 0.5 ml of ESQ Trypsin (The Mouse Kit, LEXICON) at 37°C for 15 to 20 minutes, followed by addition of ES Cell Medium (0.5 ml), to completely separate ES cell clusters into discrete cells by 35 to 40 pipettings. This cell suspension was transferred to a 15 ml centrifuge tube, and the well was washed again with 1 ml of ES Cell Medium, which was also collected in the tube. The tube was centrifuged at 1,000 rpm for 7 minutes to remove the medium. The cells were then re-suspended in 0.25 ml of ES Cell Medium and then mixed with 0.25 ml of 2 × Freezing medium. The content of the well was transferred to a cryogenic vial, frozen at -80°C and then stored in liquid nitrogen.

The cells to be used for injection into blastocysts and DNA extraction were prepared as follows. ES cell clusters were completely separated into discrete cells, one-fourth of which were provided for injection into blastocysts. One-third and two-thirds of the remaining cells were subcultured to 60 mm dishes coated with gelatin, respectively. In the former (one-third of the remaining cells), genomic DNA for Southern blotting was extracted from the cells grown to confluency, while in the latter (two-third of the remaining cells), the cells grown to confluency were aliquoted into 3 vials and frozen.

Example 3: Generation of chimeric mice from ES cells with recombinant  $\alpha$ -TTP gene

The ES cell clones, which had been confirmed to have undergone homologous recombination events, were used to form chimeric embryos by using blastocysts taken from C57BL/6J mice as host embryos. The chimeric embryos were transferred into the uterine horns of pseudopregnant mice to obtain pups. Host embryos were taken from 2-day pregnant mice by flushing their uterine tubes and uteri with Whitten's medium supplemented with 100  $\mu$ M EDTA. Eight-celled embryos or morulas were cultured in Whitten's medium for 24 hours and the resulting blastocysts were provided for injection. ES cells to be injected were subcultured for 2 or 3 days before TE treatment, and then allowed to stand at 4°C until microscopic manipulation.

As an injection pipette for ES cells, a Cook IVF polar body extrusion pipette (inner diameter: about 20  $\mu$ m) was used. A pipette for holding the embryo was prepared by longitudinally enlarging a glass microtube with an outer diameter of 1 mm (NARISHIGE) using a microelectrode-generator (model P-98/IVF, Sutter), cutting the tube at a site with an outer diameter of 50 to 100  $\mu$ m by using microforge (De Fonburun), and then reducing its bore size to 10 to 20  $\mu$ m.

Each of the injection pipettes and holding pipettes has an about 5 mm-long tip bent about 30 degrees, and was connected to a micromanipulator (LEITZ). As a

chamber for microscopic manipulation, a perforated glass slide was modified by attaching a cover slip thereto with beeswax. Two drops (about 20  $\mu$ l each) of Hepes-buffered Whitten's medium supplemented with 0.3% BSA were placed on the slide and further over-layered with liquid paraffin (Nacalai Tesque 261-37 SP). About 100 ES cells were introduced into one drop, and 10 to 15 expanded blastocysts were introduced into the other drop. 10 to 15 ES cells were injected into each blastocyst.

All microscopic manipulations were performed with an inverted microscope. The manipulated embryos were cultured for 1 to 2 hours and then transferred into the uterine horns of 2-day pseudopregnant ICR recipient females. The recipient females giving no birth past their expected date of delivery were subjected to cesarean operation and their pups were brought up by foster mothers.

When Clone L236 ES cells were injected into 40 blastocysts from C57BL/6J mice, the injection succeeded in all 40 blastocysts (100% success). These 40 blastocysts were then transferred to the uterine horns of 2-day pseudopregnant ICR recipient females, thereby obtaining 5 pups. The coat color of regions originating from homologous recombinants is a color of wild-type, while the coat color of regions originating from C57BL/6J mice is black. All the resulting 5 pups were identified as chimeric mice based on their coat color and were morphologically males. Contribution (%) of the ES cells ranged from 10% to 90%, as determined by these chimeric mice's coat color. Likewise, chimeric mice were also produced from Clone L253 ES cells. Table 1 shows scores on the generation of these chimeric mice.

**Table 1**

Scores on generation of chimeric mice

Clone No.	Injected / manipulated embryos	Transferred embryos	Nidated embryos	Pups				Coat color chimeric pups				Contribution (%) of ES cells to coat color
				Total	(%)	♂	♀	Total	(%)	♂	♀	
L228	22 / 25 (88%)	22	21 (95%)	10	45	6	4	4	40	2	2	♂(70, 70) ♀(80, 5)
L236	40 / 40 (100%)	40	24 (60%)	14	35	11	3	5	36	5	0	♂(90, 90, 90, 70, 10)

L253	75 / 77 (97%)	75	65 (87%)	11	15	7	4	8	73	5	3	♂(95, 80, 70, 50, 30) ♀(60, 50, 10)
L44	64 / 72 (89%)	64	42 (66%)	16	25	16	0	11	69	11	0	♂(90, 90, 90, 80, 80, 70, 50, 40, 30, 20, 5)

**Example 4:** Assay of homologous recombinants for germline transmission

The chimeric mice obtained in Example 3 were crossed with C57BL/6J mice to assay whether ES cell-derived pups were obtained. If the chimeric mice have germ cells derived from ES cells, the resulting pups have a wild-type coat color, whereas if they have germ cells derived from C57BL/6J mouse blastocysts, the resulting pups have a black coat color.

In the case of Clone L236 ES cells, all two male chimeric mice (No. L236-1 and -2), except for one mouse which died before sexual maturation, were found to give the germline transmission of ES cells. These two mice provided a ratio of pups of wild-type coat color to total pups of 32/44 and 19/44, respectively. In the case of Clone L254 ES cells, among 6 chimeric mice (No. L254-1 to -6), 4 mice (No. L254-1, -2, -4 and -5) were found to give the germline transmission of ES cells. These mice provided a ratio of pups of wild-type coat color to total pups of 4/15, 2/12, 2/14 and 4/9, respectively.

Next, a tail sample was taken from each pup of wild-type coat color for DNA extraction. The extracted DNA was PCR-assayed for transmission of the mutated  $\alpha$ -TTP allele, thereby confirming that the mutated  $\alpha$ -TTP allele was transmitted to both Clone L236 ES cell-derived pups and Clone L254 ES cell-derived pups.

Heterozygous deficient mice having a mutation on one allele of the  $\alpha$ -TTP gene were crossed with each other to generate homozygous deficient mice having mutations on both alleles. PCR was employed for genotype analysis of wild-type, heterozygous deficient and homozygous deficient mice. More specifically, the presence of the mutated allele was detected by PCR using the above-mentioned primer

set of ot-198 and PGK-1, while the presence of the wild-type allele was detected by PCR using a primer set of ot-198 and TTP N17 containing the sequence of Exon 1 (5' TCTCTGCAATGCCCCGCCGTGCTGTCCCG 3'; SEQ ID NO: 7). PCR using a primer set of ot-198 and TTP N17 was performed under the similar conditions as PCR using the above-mentioned primer set of ot-198 and PGK-1. The amplified products from PCR using these two primer sets were analyzed by electrophoresis. Fig. 3 shows the results obtained. Based on the electrophoresis analysis, a mouse having a detectable wild-type allele but no detectable mutated allele was identified as a wild-type mouse (Wi in Fig. 3); in contrast, a mouse having a detectable mutated allele but no detectable wild-type allele was identified as a homozygous deficient mouse (HO); and a mouse having both detectable wild-type and mutated alleles was identified as a heterozygous deficient mouse (HE).

Genotype analysis was also performed by Southern hybridization. Mouse genomic DNA was cleaved with EcoRI and electrophoresed on a 0.7% agarose gel. 15  $\mu$ g of genomic DNA was run in each lane. DNA in the gel was transferred to a Hybond N+ nylon filter (Amersham), followed by Southern hybridization. As a probe, a sequence between SmaI-SmaI-SmaI sites around Exon 1 was used (see Figs. 1 and 2). Fig. 4-1 shows the results of electrophoresis and Fig. 4-2 shows the results of Southern hybridization. As shown in Fig. 4-2, lanes 2, 3, 9, 12, 16 and 17 gave no detectable signal at a position indicated with an arrow (3.75 kbp). This suggests that mice corresponding to lanes 2, 3, 9, 12, 16 and 17 lack a sequence around Exon 1.

The above genotype analyses demonstrated that the ratio among wild-type, heterozygous deficient and homozygous deficient mice was 63:105:74 (total 242) and therefore substantially accorded with Mendel's law (1:2:1). In the case of 242 mice in total, a calculated ratio according to Mendel's law is 60.5:121:60.5.

Example 5: Analysis of  $\alpha$ -TTP expression in  $\alpha$ -TTP-deficient mice

Total RNA and protein were collected from the liver of each mouse of 6 homozygous  $\alpha$ -TTP-deficient mice, 5 heterozygous  $\alpha$ -TTP-deficient mice and 4 wild-type mice, and confirmed for expression of  $\alpha$ -TTP mRNA and protein by Northern blotting and Western blotting, respectively.

Northern blotting was performed on 10  $\mu$ g of total RNA using the entire ORF of mouse  $\alpha$ -TTP cDNA as a probe. Fig. 5-1 shows the results of Northern blotting. In addition, expression levels of  $\alpha$ -TTP mRNA were determined from PSL values of a bioimage analyzer (BAS 5000, Fuji Film), and then compared among wild-type, heterozygous and homozygous mice. Fig. 5-2 shows the results obtained. As shown in this figure, the ratio of mRNA expression levels among wild-type, heterozygous and homozygous mice was 2:1:0.

Western blotting was performed using an anti-rat  $\alpha$ -TTP polyclonal antibody and an alkaline phosphatase-conjugated anti-rabbit IgG as a secondary antibody. An AP conjugate substrate kit (Bio-Rad) was used to develop color. Fig. 6 shows the results of Western blotting. As shown in this figure, the ratio of protein expression levels among wild-type, heterozygous and homozygous mice was 2:1:0.

Each of the homozygous  $\alpha$ -TTP-deficient, heterozygous  $\alpha$ -TTP-deficient and wild-type mice was fed with  $\alpha$ -tocopherol at a concentration of 82  $\mu$ mol/kg from 4 weeks of age. All mice were then assayed for their plasma  $\alpha$ -tocopherol levels according to the method of Kim, H. S. et al. (Free Rad. Res. 28, 87-92,1998). More specifically, 50  $\mu$ l of mouse plasma collected using heparin was diluted with 950  $\mu$ l of PBS, mixed with 1 ml of 6% pyrogallol in EtOH, and then allowed to stand at 70°C for 2 minutes. After addition of 60% KOH (0.2 ml), the mixture was saponified at 70°C for 30 minutes, vortexed with 5 ml of n-hexane and 2.5 ml of H<sub>2</sub>O, and then centrifuged to collect the n-hexane layer (4 ml). The layer was evaporated and dissolved in 100 ml of ethanol, followed by HPLC analysis. HPLC was performed under the following

conditions: column: IRIKA RP18 (250 × 4 mm), mobile phase: MeOH/H<sub>2</sub>O/NaClO<sub>4</sub> (1000/2/7, v/v/w), detection: IRIKA Amperometric E-520 detector. Tocol was used as an internal standard. Fig. 7 shows the time course of  $\alpha$ -tocopherol level in each mouse. As shown in Fig. 7, the homozygous deficient mice gave a plasma  $\alpha$ -tocopherol level under the detection limit. The heterozygous deficient mice gave an about one-half plasma  $\alpha$ -tocopherol level as compared with the wild-type mice. This suggested that the expression level of  $\alpha$ -TTP was a factor in regulating blood  $\alpha$ -tocopherol level.

#### Example 6: Fetal resorption-gestation test

In view of the fact that vitamin E has been discovered as an anti-sterility agent, a fetal resorption-gestation test is widely used as a biological test for vitamin E deficiency (Biol. Syposia 12, 459, 1947). Namely, this test utilizes a phenomenon in which a pregnant animal fed with vitamin E-deficient diet does not maintain any further fetal development which leads to fetal resorption. This test was performed on the homozygous  $\alpha$ -TTP-deficient mice to biologically confirm the state of vitamin E deficiency, thereby indicating that the homozygous  $\alpha$ -TTP-deficient mice became pregnant, but showed fetal development stopped in the middle stage of pregnancy, and finally fetal resorption occurred (Table 2).

**Table 2**

Fetal resorption-gestation test in  $\alpha$ -TTP knockout mice

Genotype of female mouse	Mouse No.	×	Genotype of male mouse	Mouse No.	Nidated embryos	Regressed or resorbed fetuses	(%)*	Viable fetuses	(%)*
Homo	42	×	Homo	15	16	16	100	0	0
	44	×		17	16	16	100	0	0
	46	×		31	7	7	100	0	0
	27	×		4	0	---	---	---	---
	6	×		3	0	---	---	---	---
	Total				39	39	100	0	0
	5	×	Wild-type	C57BL/6J	8	8	100	0	0
	8	×		C57BL/6J	11	11	100	0	0
	17	×		C57BL/6J	10	10	100	0	0
	13	×		C57BL/6J	6	6	100	0	0
	Total				35	35	100	0	0

Wild-type	C57BL/6J	×	Homo	10	4	1	25	3	75
	C57BL/6J	×		2	9	2	22	7	78
	C57BL/6J	×		12	9	0	0	9	100
	C57BL/6J	×		15	9	9	100	0	0
	ICR	×		15	15	0	0	15	100
	ICR	×		15	16	2	13	14	87
	C57BL/6J	×		31	10	10	100	0	0
	ICR	×		31	16	1	6	15	94
Total					88	25	28	63	72

\* % relative to nidated embryos

In turn, when wild-type 2-celled embryos were transferred to the uterine tubes of pseudopregnant homozygous deficient mice, the mice also showed fetal development stopped during pregnancy, and finally fetal resorption occurred (Table 3).

**Table 3**

Scores on fetal resorption-gestation test in  $\alpha$ -TTP knockout homozygous females with transferred wild-type embryos

Homozygous female No	Genotype of transferred embryo	Developmental stage of transferred embryo	Transferred embryos	Nidated embryos	(%) *	Regressed or resorbed fetuses	(%) **	Viable fetuses (18 days after transfer)	(%) **
11	Wild-type (ICR)	Pronucleus	12	0	0	---	---	---	---
14	Wild-type (ICR)	Pronucleus	12	1	8%	1	100%	0	0%
7	Wild-type (ICR)	2-celled	12	11	92%	11	100%	0	0%
15	Wild-type (ICR)	2-celled	12	10	83%	10	100%	0	0%
Total			48	22	46%	22	100%	0	0%

\*: % relative to transferred embryos

\*\*: % relative to nidated embryos

These results demonstrated that vitamin E levels in pregnant females affected fetal development regardless of fetal genotype. Accordingly, the vitamin E deficient state in the homozygous  $\alpha$ -TTP-deficient mice was supported by the biological test.

### EFFECTS OF THE INVENTION

The present invention provides a mammal artificially modified to inhibit  $\alpha$ -TTP gene expression. The mammal of the present invention is very useful as a tool for understanding mechanisms for the development of diseases caused by a disrupted  $\alpha$ -TTP gene (e.g., familial isolated vitamin E deficiency) and disease conditions suspected to be associated with oxidative stress. It is also very useful as a tool for



developing a therapeutic agent and/or method for these diseases.

### BRIEF DESCRIPTION OF DRAWINGS

Fig. 1 shows a restriction site map of BS-MG6.

Fig. 2 shows the structure of a homologous recombination vector.

Fig. 3 shows results of PCR analysis using primers specific to the wild-type  $\alpha$ -TTP gene and a mutated  $\alpha$ -TTP gene, respectively.

Fig. 4-1 shows electrophoresis results of mouse genomic DNA fragments. Fig. 4-2 shows results of Southern hybridization using a probe having a sequence around Exon 1.

Fig. 5-1 shows results of Northern hybridization using mouse  $\alpha$ -TTP cDNA as a probe. Fig. 5-2 shows the intensity of signals from Northern hybridization.

Fig. 6 shows results of Western blotting using an anti-rat  $\alpha$ -TTP polyclonal antibody.

Fig. 7 shows a time course of blood tocopherol levels in mice.

## CLAIMS

1. A non-human mammal artificially modified to inhibit the expression of its endogenous gene encoding  $\alpha$ -tocopherol transfer protein.
2. The non-human mammal according to claim 1, wherein the expression of its endogenous gene encoding  $\alpha$ -tocopherol transfer protein is inhibited by disruption of at least a part of the gene or an expression regulatory region thereof.
3. The non-human mammal according to claim 1 or 2, wherein the non-human mammal is an animal belonging to Rodent.
4. The non-human mammal according to claim 3, wherein the animal belonging to Rodent is a mouse.
5. A non-human mammal cell prepared from the non-human mammal according to any one of claims 1 to 4.
6. A non-human mammal cell artificially modified to inhibit the expression of its endogenous gene encoding  $\alpha$ -tocopherol transfer protein, said cell having the ability to differentiate into an individual.
7. The non-human mammal cell according to claim 6, wherein the expression of its endogenous gene encoding  $\alpha$ -tocopherol transfer protein is inhibited by disruption of at least a part of the gene or an expression regulatory region thereof.
8. The non-human mammal cell according to claim 6 or 7, wherein the non-human mammal cell is a cell from an animal belonging to Rodent.

9. The non-human mammal cell according to claim 8, wherein the cell from an animal belonging to Rodent is a mouse cell.
10. The non-human mammal cell according to any one of claims 5 to 9, wherein the cell is an embryonic stem cell.
11. A method for producing the non-human mammal according to any one of claims 1 to 4, which comprises the steps of:
  - (a) inserting the non-human mammal cell according to any one of claims 6 to 10 into an embryo taken from a pregnant female to form a chimeric embryo, and
  - (b) transferring the chimeric embryo into the uterus of a pseudopregnant female.
12. A method for screening medicaments, which comprises using the non-human mammal according to any one of claims 1 to 4 or the non-human mammal cell according to any one of claims 5 to 10.
13. A medicament obtained by the screening method according to claim 12.

## ABSTRACT

The present invention provides a knockout animal artificially modified to inhibit  $\alpha$ -TTP gene expression. This animal is useful as a tool for understanding mechanisms for the development of familial isolated vitamin E deficiency and other diseases induced by oxidative stress (e.g., arteriosclerosis, diabetes). It is also useful as a tool for developing a therapeutic agent for these diseases.

Fig. 1




Fig. 2




Fig. 3




Fig. 4-1




Fig. 4-2




Fig. 5-1




Fig. 5-2




Fig. 6







Fig. 7



Attorney's Docket No.: 04853.0089

## DECLARATION, POWER OF ATTORNEY AND PETITION

I (We), the undersigned inventor(s), hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I (We) believe that I am (we are) the original, first, and joint (sole) inventor(s) of the subject matter which is claimed and for which a patent is sought on the invention entitled ANIMALS WITH KNOCKED OUT  $\alpha$ -TOCOPHEROL TRANSFER PROTEIN GENE

the specification of which

☐ is attached hereto.

☒ was filed on February 22, 2002 ✓ as

Application Serial No. 10/069, 251 ✓

and amended on \_\_\_\_\_

☒ was filed as PCT international application

Number PCT/JP00/ 05686 ✓

on August 24, 2000 ✓

and was amended under PCT Article 19

on \_\_\_\_\_ (if applicable).

I (We) hereby state that I (We) have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above; that I (We) do not know and do not believe that this invention was ever known or used before my invention or discovery thereof, or patented or described in any printed publication in any country before my invention or discovery thereof, or more than one year prior to this application, or in public use or on sale in the United States for more than one year prior to this application; that this invention or discovery has not been patented or made the subject of an inventor's certificate in any country foreign to the United States on an application filed by me or my legal

representatives or assigns more than twelve months before this application.

I (We) acknowledge the duty to disclose information known to be material to the patentability of this application as defined in Section 1.56 of Title 37 Code of Federal Regulations.

I (We) hereby claim foreign priority benefits under Section 119(a)-(d) of Title 35 United States Code, of any foreign application(s) for patent or inventor s certificate listed below and have also identified below any foreign application for patent or inventor s certificate having a filing date before that of the application on which priority is claimed:

Application No.	Country	Filing date	Priority claimed	
237003/1999 ✓	Japan ✓	August 24, 1999 ✓	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes	<input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes	<input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes	<input type="checkbox"/> No

I hereby claim the benefit under Section 119(e) of Title 35 United States Code, of any United States application(s) listed below.

\_\_\_\_\_  
(Application Number) (Filing Date)

\_\_\_\_\_  
(Application Number) (Filing Date)

I (We) hereby claim the benefit under Section 120 of Title 35 United States Code, of any United States application(s) listed below and, insofar as the subject

matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Section 112 of Title 35 United States Code, I (We) acknowledge the duty to disclose material information as defined in Section 1.56(a) of Title 37 Code of Federal Regulations, which occurred between the filing date of the prior application and national or PCT international filing date of this application:

Application Serial No.	Filing Date	Status (pending, patented, abandoned)
_____	_____	_____
_____	_____	_____
_____	_____	_____

And I (We) hereby appoint: FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER, L.L.P. CUSTOMER NUMBER 22,852

I(We) hereby request that all correspondence regarding this application be sent to the firm of FINNEGAN, HENDERSON, FARABOW, GARRETT & DUNNER, L.L.P., whose Post office address is: 1300 I Street, N.W., WASHINGTON, D.C. 20005 U.S.A.

I (We) declare further that all statements made herein of my (our) knowledge are true and that all statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

1-00 Keizo INOUE  
NAME OF FIRST SOLE INVENTOR

Keizo Inoue  
Signature of Inventor

May 8, 2002  
Date

Residence: Tokyo, Japan JPX

Citizen of: Japan ✓

Post Office Address: 17-605,  
Kantozaimukyoku-Ecchujimajyutaku,  
1-3, Ecchujima, Koto-ku, Tokyo  
135-0044 Japan

2-00 Hiroyuki ARAI  
NAME OF SECOND JOINT INVENTOR

Hiroyuki Arai  
Signature of Inventor

May 8, 2002  
Date

Residence: Tokyo, Japan JPX

Citizen of: Japan ✓

Post Office Address: 5-35-8-604,  
Koishikawa, Bunkyo-ku, Tokyo  
112-0002 Japan

3-00 Makoto ARITA  
NAME OF THIRD JOINT INVENTOR

Makoto Arita  
Signature of Inventor

May 8, 2002  
Date

Residence: Kanagawa, Japan JPX

Citizen of: Japan ✓

Post Office Address: 1-21-4-206,  
Azamino, Aoba-ku, Yokohama-shi,  
Kanagawa 225-0011 Japan

4-00

Kou-ichi JISHAGE

NAME OF FOURTH JOINT INVENTOR

Kouichi Jishage

Signature of Inventor

May 8, 2002

Date

Residence: Shizuoka, Japan JPXCitizen of: Japan ✓Post Office Address: 2-3-3,  
Sanomiharashidai, Mishima-shi,  
Shizuoka 411-0041 Japan

5-00

Hiroshi SUZUKI

NAME OF FIFTH JOINT INVENTOR

Hiroshi Suzuki

Signature of Inventor

May 8, 2002

Date

Residence: Shizuoka, Japan JPXCitizen of: Japan ✓Post Office Address: 1150-22,  
Shimotogari, Nagaizumi-cho,  
Sunto-gun, Shizuoka 411-0943 Japan

SEQUENCE LISTING

<110> Keizo Inoue

Hiroyuki Arai

<120> An animal with a knocked out  $\alpha$ -tocopherol transfer protein gene

<130> PH-1014-PCT

<150> JP P1999-237003

<151> 1999-08-24

<160> 7

<170> PatentIn Ver. 2.0

<210> 1

<211> 837

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (1).. (834)

<400> 1

atg gca gag atg cgg ccg ggg cca ttg gtt ggg aaa cag ctc aac gag 48

Met Ala Glu Met Arg Pro Gly Pro Leu Val Gly Lys Gln Leu Asn Glu

1

5

10

15

ctg ccc gac cac tcg ccg ctg ctc cag ccc ggc ctg gcg gag ctc agg 96  
 Leu Pro Asp His Ser Pro Leu Leu Gln Pro Gly Leu Ala Glu Leu Arg  
                   20                  25                  30

cgc cgg gtg cag gag gca ggc gtc ccg cag acc ccg cag cct ctc aca 144  
 Arg Arg Val Gln Glu Ala Gly Val Pro Gln Thr Pro Gln Pro Leu Thr  
                   35                  40                  45

gac gct ttc ctg ctg cgc ttc ctg cgc gcc cgg gat ttc gat ctg gat 192  
 Asp Ala Phe Leu Leu Arg Phe Leu Arg Ala Arg Asp Phe Asp Leu Asp  
                   50                  55                  60

ctg gcc tgg cgc tta atg aaa aac tat tat aaa tgg cga gca gaa tgc 240  
 Leu Ala Trp Arg Leu Met Lys Asn Tyr Tyr Lys Trp Arg Ala Glu Cys  
                   65                  70                  75                  80

cca gaa tta agt gca gat cta cgc cct aga agt atc ctt gga ctt ctg 288  
 Pro Glu Leu Ser Ala Asp Leu Arg Pro Arg Ser Ile Leu Gly Leu Leu  
                   85                  90                  95

aaa gct ggc tac cat ggc gtg ctc agg tcc cgg gat tct act ggc agt 336  
 Lys Ala Gly Tyr His Gly Val Leu Arg Ser Arg Asp Ser Thr Gly Ser  
                   100                  105                  110

aga gtt ctc att tac aga att gca tac tgg gac cca aaa gtt ttt aca 384  
 Arg Val Leu Ile Tyr Arg Ile Ala Tyr Trp Asp Pro Lys Val Phe Thr  
                   115                  120                  125

gct tat gat gta ttt cgt gta agt ctg atc aca tca gag ctc att gta 432  
 Ala Tyr Asp Val Phe Arg Val Ser Leu Ile Thr Ser Glu Leu Ile Val



130	135	140	
cag gag gtg gaa act caa cgc aat gga gtt aaa gct ata ttt gac ctg	480		
Gln Glu Val Glu Thr Gln Arg Asn Gly Val Lys Ala Ile Phe Asp Leu			
145	150	155	160
gaa ggc tgg cag gtt tct cat gct ttc caa att acc cca tct gta gcc	528		
Glu Gly Trp Gln Val Ser His Ala Phe Gln Ile Thr Pro Ser Val Ala			
	165	170	175
aag aag att gct gct gta ctt aca gat tcc ttt cca ctg aaa gtt cgt	576		
Lys Lys Ile Ala Ala Val Leu Thr Asp Ser Phe Pro Leu Lys Val Arg			
	180	185	190
ggg atc cat ttg ata aat gag cca gtc att ttc cat gct gtc ttc tcc	624		
Gly Ile His Leu Ile Asn Glu Pro Val Ile Phe His Ala Val Phe Ser			
	195	200	205
atg att aaa cca ttt ctg act gaa aag att aag gac cgg att cat ctg	672		
Met Ile Lys Pro Phe Leu Thr Glu Lys Ile Lys Asp Arg Ile His Leu			
	210	215	220
cac ggg aac aac tac aaa tca agc atg ctt cag cac ttc cca gac att	720		
His Gly Asn Asn Tyr Lys Ser Ser Met Leu Gln His Phe Pro Asp Ile			
	225	230	235
ctt cct cgg gaa tat ggc ggt aaa gag ttc tcc atg gag gat att tgt	768		
Leu Pro Arg Glu Tyr Gly Gly Lys Glu Phe Ser Met Glu Asp Ile Cys			
	245	250	255

cag gag tgg aca aat ttt ata atg aag tct gaa gat tat ctc agc agc 816  
 Gln Glu Trp Thr Asn Phe Ile Met Lys Ser Glu Asp Tyr Leu Ser Ser  
 260 265 270

att tct gag acc atc caa tga 837  
 Ile Ser Glu Thr Ile Gln  
 275

<210> 2

<211> 278

<212> PRT

<213> Mus musculus

<400> 2

Met Ala Glu Met Arg Pro Gly Pro Leu Val Gly Lys Gln Leu Asn Glu  
 1 5 10 15

Leu Pro Asp His Ser Pro Leu Leu Gln Pro Gly Leu Ala Glu Leu Arg  
 20 25 30

Arg Arg Val Gln Glu Ala Gly Val Pro Gln Thr Pro Gln Pro Leu Thr  
 35 40 45

Asp Ala Phe Leu Leu Arg Phe Leu Arg Ala Arg Asp Phe Asp Leu Asp  
 50 55 60

Leu Ala Trp Arg Leu Met Lys Asn Tyr Tyr Lys Trp Arg Ala Glu Cys  
 65 70 75 80

Pro Glu Leu Ser Ala Asp Leu Arg Pro Arg Ser Ile Leu Gly Leu Leu  
                             85                            90                            95

Lys Ala Gly Tyr His Gly Val Leu Arg Ser Arg Asp Ser Thr Gly Ser  
                             100                            105                            110

Arg Val Leu Ile Tyr Arg Ile Ala Tyr Trp Asp Pro Lys Val Phe Thr  
                             115                            120                            125

Ala Tyr Asp Val Phe Arg Val Ser Leu Ile Thr Ser Glu Leu Ile Val  
                             130                            135                            140

Gln Glu Val Glu Thr Gln Arg Asn Gly Val Lys Ala Ile Phe Asp Leu  
                             145                            150                            155                            160

Glu Gly Trp Gln Val Ser His Ala Phe Gln Ile Thr Pro Ser Val Ala  
                             165                            170                            175

Lys Lys Ile Ala Ala Val Leu Thr Asp Ser Phe Pro Leu Lys Val Arg  
                             180                            185                            190

Gly Ile His Leu Ile Asn Glu Pro Val Ile Phe His Ala Val Phe Ser  
                             195                            200                            205

Met Ile Lys Pro Phe Leu Thr Glu Lys Ile Lys Asp Arg Ile His Leu  
                             210                            215                            220

His Gly Asn Asn Tyr Lys Ser Ser Met Leu Gln His Phe Pro Asp Ile  
                             225                            230                            235                            240

Leu Pro Arg Glu Tyr Gly Gly Lys Glu Phe Ser Met Glu Asp Ile Cys  
                   245                  250                  255

Gln Glu Trp Thr Asn Phe Ile Met Lys Ser Glu Asp Tyr Leu Ser Ser  
                   260                  265                  270

Ile Ser Glu Thr Ile Gln  
                   275

<210> 3

<211> 2748

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (1139).. (1342)

<400> 3

gaattcaaag cctcagccc ggtaaccaag caccacagcc agctctcttt gtgattcagg 60

ggttcacacc acaacacagc cgtttggcct tgttccctgg tgtttgctta atgttctcct 120

acaccatgga ggagatttac cctcgtcctt ttacttcca gccacacaa aaatgaaaaa 180

cgctccaag gcaagagttc tgttttgagg atatcctcaa taatcggaac atggctctta 240

cccaagagcc actccatcag acattcttgc tctgagttcc ttttaaggcct ctttactcgc 300

caaaatcagt gttttgtgaa catgcactgc atattaagag gagttagttt tgggacttt 360  
 ctctgttca ggtggcagtt caagtgtagg ataattttaa tggaaatgaa ggaaaaatac 420  
 ttccgtgtgt tcatcagat ttccgggtca tctctgtgta ttcttcagca gacatccttc 480  
 aggttcctta agtaagggtt ttgattgaga gactgggtgc atctaaacac atacatcgtt 540  
 agtgittaaa aatgtgacct cccccacccg ctctcctttc tctagtagag ccagatgcca 600  
 agatctggaa gcattttcct ggagagaagc aaggaggagg aggaggagac tgccaaaagg 660  
 tgacttcctt gagttacatt ttggaaacta gttagaatgc cagagatggc ctgagctcag 720  
 ccttaaggaa ggggtcagga ggagggttcc tgagtgtctg ctaccaagc taattaaaga 780  
 gccgtttaca gtgttccctg aticccaaaac ggacagaggg ggaagggcaa cgaggaaagg 840  
 gtgagaaaag tctctggcag cctgattata aacatcccaa gtaacttttc gacttcccgt 900  
 tctttaggtt caacactagt gactttccct tccccggga ctggctgcgg ttaccttggt 960  
 gagcaccgga gggcaccacg tgggttctt taagagggcg ccgtgaccct tgcaccggcg 1020  
 gggcacggga gatcggggcg gcccgggtga gtgtgcgtgg ggcggcgtcc acggcggggg 1080  
 gcggagggtg gctctgggcc cgcacttttc cccctgtcgc cgggacagca cggcgggc 1138  
 atg gca gag atg cgg ccg ggg cca ttg gtt ggg aaa cag ctc aac gag 1186  
 Met Ala Glu Met Arg Pro Gly Pro Leu Val Gly Lys Gln Leu Asn Glu

1	5	10	15	
ctg ccc gac cac tcg ccg ctg ctc cag ccc ggc ctg gct gag ctc agg	1234			
Leu Pro Asp His Ser Pro Leu Leu Gln Pro Gly Leu Ala Glu Leu Arg				
20	25	30		
cgc cgg gtg cag gag gca ggc gtc ccg cag acc ccg cag cct ctc aca	1282			
Arg Arg Val Gln Glu Ala Gly Val Pro Gln Thr Pro Gln Pro Leu Thr				
35	40	45		
gac gct ttc ctg ctg cgc ttc ctg cgc gcc cgg gat ttc gat ctg gat	1330			
Asp Ala Phe Leu Leu Arg Phe Leu Arg Ala Arg Asp Phe Asp Leu Asp				
50	55	60		
ctg gcc tgg cgc gtaagtgtgc accgggggcg ggcagagctc ggcgacggcg	1382			
Leu Ala Trp Arg				
65				
gaatccacgc gcgccgagcg tggcagtgtg actgcaggcg cgcccagaac cccgatttcg	1442			
ccccgccga tgttttggtc cccgccgccg cgaggacatc ccttggaacta ctagggctct	1502			
tgggaattaa acaaagtgga gatccctgtc ccccggggtg ctcagctgtg ttaactgaat	1562			
agataactag gtgtggacag aggacgacga aatggacatc taaaggcatc ttgaaaaaga	1622			
ctatgttaat agagctaaat gcacagtgtg gcatgtttga macccagggc agtacagatg	1682			
atttctttta tgtttcaggt attcacaaca cactggcctt ggggcaagag agatggggcc	1742			

ttagggtcag ggagatgcga ccttgacttt gtccctcttg gggtcagcac ccttatctgt 1802  
 tcagtaactg tgaggacatg acagtagitt cgagaattgc acattaacct ggaatgctag 1862  
 aacaagatgt gccaaacct gtgcttggca cggagaaagt agtcagtgt cagcaggctg 1922  
 cggatttcca acatgccctg ggtttatgaa acttttttta ttggataagc accaagtatg 1982  
 gcaaaaaaca ccacaaacaa tacaaaacag gaaaaacdtc aaaggaattt cctaaaagaa 2042  
 aagaatticc caacacaaac tctagttaga ccttgaggac ccagaagtat ggcattacct 2102  
 gttacgtcaa gccigtglaa caatgtcacg caaacatgcg ctgtgagttt atttttcctt 2162  
 tgcaaatctc aactgcatgc tgttatagaa tcaggtcatg tgaacatgtg ctacaccta 2222  
 ctactctttt gggaatatct agtcagtttt ttgtttgttg ctgtagagat tgttaccggg 2282  
 cgggtgctgt taggccctat gctgatcggt catccctaca ttcagtgtat ggggaccag 2342  
 cgctgccatg ttactgttc atctcttcat ttcatltgga gtttctcctt ctttttcttt 2402  
 ctttttcttt tcttttcttt ttaataacca cacactgcct agcagtatac aaatgccatc 2462  
 aacaggltgag tattttcttc tctccctgac tgcattctaa ttggtctttg tctgtacaca 2522  
 taaattggaa catatcctta ttgaacaaat ccatcagttg ctgaagcacg acgcagacat 2582  
 gtttactgtt gaggagcgca ccaccttgc agggagtttt cagtgtttgc tactctgatg 2642

aaatgcacac tgcatagtga cgtctttttc tctctattgt tatgtacact gtctttacaa 2702

atrggatgta tgccctgctag atgaggatag ttttgcattt cattat 2748

<210> 4

<211> 68

<212> PRT

<213> Mus musculus

<400> 4

Met Ala Glu Met Arg Pro Gly Pro Leu Val Gly Lys Gln Leu Asn Glu

1

5

10

15

Leu Pro Asp His Ser Pro Leu Leu Gln Pro Gly Leu Ala Glu Leu Arg

20

25

30

Arg Arg Val Gln Glu Ala Gly Val Pro Gln Thr Pro Gln Pro Leu Thr

35

40

45

Asp Ala Phe Leu Leu Arg Phe Leu Arg Ala Arg Asp Phe Asp Leu Asp

50

55

60

Leu Ala Trp Arg

65

<210> 5

<211> 28

<212> DNA

<213> Artificial Sequence



<400> 5

gctaaagcgc atgctccaga ctgccttg

28

<210> 6

<211> 31

<212> DNA

<213> Artificial Sequence

<400> 6

agccccacaca aaaatgaaaa acgtctccaa g

31

<210> 7

<211> 28

<212> DNA

<213> Artificial Sequence

<400> 7

tctctgcaat gcccgccgtg ctgtcccg

28

<210> 8

<211> 22

<212> DNA

<213> Artificial Sequence

<400> 8

aggaattcat ggcagagatg cg

22

<210> 9

<211> 22

